DEPARTMENT ACTIVITIES

Pharm D faculties and students attended 71 st Indian Pharmaceutical Congress at Chennai on the Theme Healthcare system -Role of regulators, Pharmavision 2030 on 20,21 December 2019.

Pharm D students conducted National Pollution Prevention Day on 6 January, 2020





DIC ACTIVITIES	NUMBER	Answers for Quiz					
			8.ð	J.₽	3.D	A.S	A.f






A Newsletter on

An Update on Clinical Research and Drug Information

Volume : 6	Issue : 1		January -April 2020			
An Official Publication from Department of Pharmacy Practice, Swamy Vivekanandha College of Pharm Elayampalayam, Tiruchengode - 637 205, Namakkal (Dt), Tamilnadu, Phone : 04288-234417, E-mail : svcpdpic@gmail.com	acy,	Patron Advisory Board Chief Editor Editorial Board	 Prof. Dr. M. Karunanithi Dr. S. Arthanareeswaran Dr. K.SreeraaghanidhiArthanareeswaran Dr. G. Murugananthan Dr. M. Rangapriya Mr. S. Anandkumar, Mrs. P. Parkavi Rani Dr.S. Harini, Ms.Kalaivani 			
PHARMACIST DESK						
Creutzfeldt-Jakob disease		corneal grafts, dural grafts, or electrode implar human growth hormone.				
One state to be the balance of the second state of the second sec	La subtra					

Creutzfeldt–Jakob disease also known as classic Creutzfeldt-Jakob disease, is a fatal degenerative brain disorder. CJD is caused by a protein known as a prion. Infectious prions are misfolded proteins that can cause normally folded proteins to become misfolded. Causes



The cause of Creutzfeldt-Jakob disease and other TSEs appears to be abnormal versions of a kind of protein called a prion. Normally these proteins are harmless. But when they're misshapen, they become infectious and can harm normal biological processes. SIGNS AND SYMPTOMS

Personality changes, Anxiety, Depression, Memory loss, Impaired thinking, Blurred vision or blindness, Insomnia, Difficulty speaking, Sudden, jerky movements, Coma

Transmission

The defective protein can be transmitted by contaminated harvested human brain products,

Drug and Poison Information Queries Please Contact : Email - svcpdpic@gmail.com, Phone : 04288 - 234417.



its and

It can be familial in which a mutation has occurred in the gene for PrP, PRNP.

Types

- * Sporadic caused by the spontaneous misfolding of prion-protein in an individual.
- * Familial caused by an inherited mutation in the prionprotein gene.
- Acquired CJD caused by contamination with tissue from an infected person.

TREATMENT

There was no cure for CJD. Some of the symptoms like twitching can be managed, but otherwise treatment is palliative care. Psychiatric symptoms like anxiety and depression can be treated with sedatives and antidepressants. Myoclonic jerks can be handled with clonazepam or sodium valproate. Opiates can help in pain. Seizures are very uncommon and can be treated with antiepileptic drugs.

PREVENTION

Destruction of surgical instruments used on the brain or nervous tissue of someone with known or suspected CJD.Single-use kits for spinal taps. Tight restrictions on importation of cattle from countries where BSE is common.

Restrictions on animal feed.

Strict procedures for dealing with sick animals.

CASE REPORT

Traumatic brain injury

A2 years old male child was admitted with the complaints of swelling and pain in head due to falling from bed, changes in sleep habit, persistent headache. The child had no past medical and medication history. The vital shows that elevated blood pressure (130/90mmhg) and other parameters were found to be normal. The child was diagnosed with traumatic injury on brain with soft tissue injury. The child was initiated on syrup .ossopan 3ml twice daily, syrup. dilantin 5ml twice daily, syrup. piracetam 2.5 ml twice daily, tab. bizlo 20 mg twice daily for four days.

Conclusion

Traumatic brain injury (TBI) is a non-degenerative, non congenital force to the brain from external mechanical force. A computed tomography scan (CT or CAT scan) is the gold standard for the radiological assessment of a Traumatic brain injury. TBI is temporary or permanent according to the severity of injury. It can be cured by TBI rehabilitation. TBI Rehabilitation aims to reduce their pain, promote healing, improving their cognitive functions, voluntary moments and increase their quality of life.

Ref :

Ms. JANANI PRRIYA .S, Pharm .D, Intern,

	QUIZ					
1. Which one of this is a diagnostic agent for myasthenia gravis?						
A. Endrophonium	B. Physositgmine					
C. Neostigmine	D. Pyridostigmine					
2. lodohydroxyquinoline causes:						
A. Jaundice	B. Eye defects					
C. Diarrhoea	D. Ataxia					
3. Which one(s) of the following a	ntihypertensive medications may be associated					
with depressive syndrome develo	pment?					
A. Clonidine	B. Guanethidine					
C. Methyldopa	D. All of the above					
-	pomorphine and ondansetron may result in which					
one(s) of the following adverse ef	fect/effects?					
A. Loss of consciousness	B. Significant hypotension					
C. Both	D. Neither					
5. Example of an "indirect acting" adrenergic agonist:						
A. Phenylephrine	B. Cocaine					
C. Oxymetazoline	D. Isoproterenol					

Ref :

IRENE GEORGE, Pharm D., Intern,

	RECENTLY APPROVED DRUGS BY FDA						
S. No.	DRUG NAME	DOSE	DOSAGE	INDICATIONS	APPROVED ON		
1.	Tepezza	500 mg	Injection	Thyroid eye disease	21.01.2020		
2.	Numbrino nasal solution	40 -160 mg	Nasal solution	Nasal anesthesia	10.01.2020		
3.	Ayvakit	300 mg	Tablets	Gastrointestinal Stromal Tumor	09.01.2020		
4.	Ubrelvy	50-100 mg	Tablets	Migraine	23.12.2019		
5.	Dayvigo	10 mg	Tablets	Insomnia	20.12.2019		

Ref: www.fda.com

NEW DRUG PROFILE

OXBRYT	Ά	XCOPRI			
Generic name : Voxe	lotor	Genericname	: Cenobamate		
Drug class : Hem	noglobin Oxygen – Affinity	Drug class	: Anti-epileptic Drug		
Modu	ulators	Dosage forms and strength	: 400 mg tabelet		
Dosage form and strength : 500 m	ng tablet	Indication	: Partial-onset Seizures		
Indication : Sickl	le cell disease	МоА	: To reduce repetitive		
Moa : It is a	a hemoglobin S		neuronal firing by inhibiting voltage –gated sodium		
polyn	nerization inhibitor that		currents. It is also a		
binds	s to HbS with a 1:1		positive allosteric		
stoich	hemistry and exhibits		modulator of the		
prefe	erential partitioning to		γ- aminobutyric acid ion		
RBC	S		channel.		
ADR : Hype	ersensitivity, Fatigue,	ADR	: QT Shortening,Suicidal Behaviour, Somnolenence		
Pyrex	xia, Nausea, Headache.		and Fatigue		
Storage : store	at or below 30°C	Storage	: Store at 20-25°C		
Drug Approved on : 25.11	1.2019	Drug Approved on	: 21.11.2019		

Ref : IRINE JACOB, Pharm D, Intern,